

AD-A274 745



AD _____

CONTRACT NO: DAMD17-90-C-0106

TITLE: PATHOBIOLOGY OF HIV IN THE HUMAN MONOCYTE-MACROPHAGE

PRINCIPAL INVESTIGATOR: Jerome E. Groopman, M.D.

CONTRACTING ORGANIZATION: New England Deaconess Hospital
185 Pilgrim Road
Boston, Massachusetts 02215

REPORT DATE: December 10, 1993

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and
Development Command, Fort Detrick
Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC
S ELECTE D
JAN 14 1994
C

94-1 13 015

388 283 94-01520

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION			1b. RESTRICTIVE MARKINGS			
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE						
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)			
6a. NAME OF PERFORMING ORGANIZATION New England Deaconess Hospital		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION			
6c. ADDRESS (City, State, and ZIP Code) 185 Pilgrim Road Boston, MA 02215			7b. ADDRESS (City, State, and ZIP Code)			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract No. DAMD17-90-C-0106			
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21702-5012			10. SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO. 63105A	PROJECT NO. 3M2 63105870.	TASK NO. AA	WORK UNIT ACCESSION NO. 019
11. TITLE (Include Security Classification) Pathobiology of HIV in the Human Monocyte-Macrophage						
12. PERSONAL AUTHOR(S) Groopman, Jerome E.						
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 90/09/28 TO 93/09/27		14. DATE OF REPORT (Year, Month, Day) 93/12/10		
15. PAGE COUNT 20						
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD	GROUP	SUB-GROUP				
06	03		HIV, Monocytes, Host Defense			
06	13		RA I, AIDS, Viral Tropism, Cytokine			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We have studied various aspects of HIV biology within the monocyte-macrophage. These studies included interactions of newly discovered cytokines in promoting monocytic cell development in the context of HIV infection. Both interleukin-3 and kit ligand/stem cell factor were found to augment myelopoiesis in vitro and protect monocyte-macrophage from toxic effects of Zidovudine and ganciclovir. Kit ligand/stem cell factor had no upregulatory effect on HIV transcription while interleukin-3 did. These studies set the stage for the use of these cytokines to reduce the cytotoxicity of important agents such as Zidovudine and ganciclovir in the treatment of HIV disease and its complicating opportunistic infections. We also discovered that the long-terminal repeat (LTR) of HIV contains motifs responsive to steroid hormone receptors. There appears to be a complex interplay among these transcription factors in terms of modulating virus expression in the monocyte-macrophages. These studies open a new avenue of research with regard to therapeutics based on retinoic acids. Efforts were initiated to introduce synthetic genes capable of inhibiting HIV into the monocyte-macrophages in novel adenoassociated virus (AAV) vectors. We have successfully constructed prototype vectors using the AAV backbone and found a high transduction efficiency in macrophages. Future studies will examine the optimal constructs to inhibit HIV replication in these cells						
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified			
22a. NAME OF RESPONSIBLE INDIVIDUAL Mrs. Virginia Miller			22b. TELEPHONE (Include Area Code) 301-619-7328		22c. OFFICE SYMBOL SGRD-RMI-S	

DTIC QUALITY INSPECTED 8

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____Where copyrighted material is quoted, permission has been obtained to use such material.

____Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

____In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

~~SA~~ For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45 CFR 46.

OR In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

PI Signature *James P. [unclear]* Date *12/1/93*

Pathobiology of HIV in the Human Monocyte-Macrophage

TABLE OF CONTENTS

ELEMENTS	PAGE
1. FRONT COVER	1
2. DD FORM 1473	2
3. FOREWORD	3
4. TABLE OF CONTENTS	4
FINAL REPORT	
5. INTRODUCTION	5
6. BODY	5
7. CONCLUSIONS	8
8. REFERENCES	10
9. APPENDIX	11

INTRODUCTION

Our research program explores the interactions of HIV with monocyte-macrophages. Monocyte-macrophages are important host defense cells which both interact with invading pathogens and present antigens to other immune effector cells. HIV is able to infect monocyte-macrophages via the CD4 cell surface structure and both replicate within these cells as well as potentially alter their function (1-3). The purpose of the present work is to determine the regulation of HIV within this cell type, the consequences of infection particularly with regard to cytokine generation, and to ultimately use this information for the clinical benefit of individuals with HIV disease.

BODY

A. Cytokine Biology

We pursued the identification of a novel cytokine termed "kit ligand/stem cell factor" which is produced by mesenchymal cells including bone marrow stromal fibroblasts (4,5). Because this cytokine was considered as a potential therapeutic in stimulating stem cell proliferation for purposes of immune reconstitution, we addressed whether the kit ligand/stem cell factor might alter HIV replication in monocyte-macrophages. Standard cultures of monocyte-macrophages were established in vitro using tropic isolates of HIV including BAL and 9533. We found no upregulation or effect on HIV within the monocyte-macrophage following exposure of the cells to kit ligand/stem cell factor.

This suggested to us that this cytokine, as well as other early acting hematopoietic growth factors, might of therapeutic use in people with HIV. To that end, we studied myeloid progenitor development (CFU-GM) and erythroid progenitor development (BFU-E) in vitro in the presence of important therapeutics for people with AIDS including AZT, ganciclovir as well as the inflammatory cytokine tumor necrosis factor alpha (TNF- α) (Table 1) (Figure 1).

Final Report

(Period 09/28/90 - 09/27/93)

Contract No: DAMD17-90-C-0106

We had previously determined, as had other groups, that TNF- α could upregulate HIV expression in monocyte-macrophages as well as T cells (6,7). Comparative in vitro studies in the laboratory examined the effects of recombinant human interleukin-3 with those effects from recombinant kit ligand/stem cell factor. We found that there were protective effects against both myelotoxic drugs and inflammatory cytokines such as TNF- α utilizing interleukin-3, with a discernable but less potent protective effect seen with kit ligand/stem cell factor. We confirmed that interleukin-3 may moderately upregulate HIV replication in monocyte-macrophages. Using these data, we established a framework upon which certain cytokines such as the kit ligand/stem cell factor may be used to promote the development of myeloid cells, including monocyte-macrophages, in the presence of HIV infection without stimulating replication of the retrovirus.

B. Cytokine Signalling Pathways in Monocyte-Macrophages

Based on our prior studies of TNF- α effects on HIV in monocyte-macrophages, we sought to further explore the pathways of signal transduction in myeloid cells in the context of this inflammatory cytokine. The purposes of these studies was to better understand those biochemical mediators in the monocyte-macrophage which could contribute to upregulation of HIV expression and thereby to seek novel antiviral interventions. We focused on the sphingomyelin pathway, specifically the intermediate ceramide (8). We found that ceramide was a potent upregulator of HIV replication, particularly using monocytic cell lines such as U1 (a subclone of U937) in which viral transcription is minimal. We found that exogenous ceramide was a potent upregulator of HIV expression in monocytic cells, providing direct evidence for signal transduction intermediates in the TNF-sphingomyelin pathway. Potential inhibitors of this pathway might be considered as ultimate therapeutic agents for HIV disease (Figure 2 & 3).

C. Steroid Hormone Receptors

A major part of our program has involved the study of transcriptional regulation via the negative regulatory element

Final Report

(Period 09/28/90 - 09/27/93)

Contract No: DAMD17-90-C-0106

(NRE) of the HIV long-terminal repeat (LTR). The NRE has motifs which could interact with the family of transcription factors that act as steroid hormone receptors. We first found that a newly identified steroid hormone receptor termed ARP tightly bound to a motif in the NRE. We also found that there was a complex interaction among the many members of this steroid hormone receptor family with regard to homodimer and heterodimer formation. Most potent with regard to binding to the responsive motifs in the HIV negative regulatory element were the steroid hormone receptors for trans-retinoic acid and 9-cis-retinoic acid (RAR- α and RXR- α) (9) (Figure 4).

We recently found that there was an excellent correlation with regard to downregulation of HIV expression in different monocyte-macrophage cell lines and in different T cell lines with regard to level of expression of these steroid hormone receptors in the cells. This correlation provides indirect but important data that retinoids may be useful in modulating HIV replication in different cell types (Figure 5).

D. Transcription Regulation of HIV, Independent of the NF-Kappa B Structure

We found that certain cytokines could cause transcriptional activation by the HIV LTR independent of the NF-kappa B structure. This was most dramatically seen in viruses with deletion point mutations in the two NF-kappa B sequences present in the HIV LTR. A primitive cell line with myeloid and megakaryocytic properties called CMK was studied in this regard. Induction of differentiation with the phorbol diester PMA led to transactivation despite the absence of intact NF-kappa B sequences. In the context of the pathobiology of HIV in the monocyte-macrophage, we then focused on GM-CSF treatment of THP-1 monocytic cells transfected with these NF-kappa B mutants. There was excellent transactivation of these cells as well. Sequential deletions in the LTR demonstrated the minimal sequences required for transactivation which are clustered around the TATA box of the HIV LTR.

E. Transduction of Monocyte-Macrophages with Adenoassociated Virus Vectors (AAV)

In order to dissect the effects of different gene products within the monocyte-macrophage, as well as use artificial genes which might inhibit HIV, we sought to identify a vector system which had a high transduction efficiency for resting cells such as macrophages. We have been fortunate in obtaining adenoassociated virus (AAV) vector and have conducted a series of studies to determine ability of these vectors to transduce monocyte-macrophages. Our initial studies are very promising. Using specific marker genes including CAT as well as Beta-Gal, we have found excellent transduction efficiency (as high as 50%) in some experiments. No cytopathic effects of the AAV vector have been discernable in the monocyte-macrophage.

We will now proceed by introducing specific HIV gene products, including VPR and VPU, into the monocyte-macrophage under the control of various promoters. The effects of the gene products on monocyte-macrophage function with regard to antigen presentation in cytokine generation will be studied. In addition, we will introduce antisense constructs for purposes of inhibiting HIV within the monocyte-macrophage by exploiting the AAV vector.

CONCLUSION

The program has succeeded in addressing many of the goals set out three years ago. We have an improved understanding of signal transduction pathways of important cytokines such as TNF- α within the monocyte-macrophage. We have identified the novel cytokine kit ligand/stem cell factor which does not upregulate HIV within the monocyte-macrophage but leads to enhanced progenitor development, thereby providing an opportunity for potential stem cell expansion and immune reconstitution. A new mechanism of transcriptional activation of HIV in the monocyte-macrophage via steroid hormone receptor interaction with motifs in the HIV LTR was identified. Manipulation of different retinoids and other steroids could lead to development of novel therapeutics. Finally, the first steps in

Final Report

(Period 09/28/90 - 09/27/93)

Contract No: DAMD17-90-C-0106

gene therapy targeting resting cells such as macrophages were taken with a new class of vectors, AAV, with very positive results to date.

The work supported by this contract thereby contributed to both the basic understanding of the pathobiology of HIV in the monocyte-macrophage as well as potential new therapeutic interventions for people with HIV disease.

Final Report

(Period 09/28/90 - 09/27/93)

Contract No: DAMD17-90-C-0106

REFERENCES

1. Gartner S, Markovits P, Markovits DM, Kaplan MH, Gallo RC. The role of mononuclear phagocytes in HTLV-III/LAV infection. *Science*. 1986; 223:215-219.
2. Hammer SM, Gillis JM, Groopman JE, Rose RM. In vitro modification of human immunodeficiency virus infection by granulocyte-macrophage colony-stimulating factor and gamma interferon. *Proc Natl Acad Sci USA*. 1986; 83:8734-8738.
3. Salahuddin SZ, Rose RM, Groopman JE, Markham PD, Gallo RC. Human T-lymphotropic virus type III infection of human alveolar macrophages. *Blood*. 1986; 68:281-284.
4. Avraham H, Scadden DT, Chi S, Broudy VC, Zsebo KM, Groopman JE. Interaction of human bone marrow fibroblasts with megakaryocytes: Role of the c-kit ligand. *Blood*. 1992; 80:1679-1684.
5. Avraham H, Vannier E, Cowley S, Jiang S, Chi S, Dinarello C, Zsebo K, Groopman JE. Effects of stem cell factor, c-kit ligand, on human megakaryocytic cells. *Blood*. 1992; 79:365-371.
6. Butera ST, Roberts BD, Folks TM. Regulation of HIV-1 expression by cytokine networks in a CD4+ model of chronic infection. *J Immunol*. 1993; 150:625-634.
7. Folks TM, Clouse KA, Justement J, Rabson A, Duh E, Kehrl JH, Fauci AS. Tumor necrosis factor- α induces expression of human immunodeficiency virus in a chronically infected T cell clone. *Proc Natl Acad Sci USA*. 1989; 86:2365-2368.
8. Papp B, Zhang D, Groopman JE, Byrn RA. Stimulation of human immunodeficiency virus type 1 expression by ceramide. Submitted to *AIDS Research and Human Retroviruses*, 1993.
9. Yamaguchi K, Groopman JE, Byrn RA. The regulation of HIV by retinoic acid correlates with cellular expression of the retinoic acid receptors. Submitted to *AIDS*, 1993.

Final Report
(Period 09/28/90 - 09/27/93)
Contract No: DAMD17-90-C-0106

APPENDIX

Table 1

Figure 1 through 5

Final Report

(Period 09/28/90 - 09/27/93)

Contract No: DAMD17-90-C-0106

FIGURE LEGEND

Figure 1: BFU-E (left column) or CFU-GM (right column) in the presence of ZDV (0, 0.05, 0.1 and 1.0 μ M), GAN (0, 0.1, 0.2, and 1.0 μ M), IFN (0, 100, 1000 and 5000 U/ml), TGF (0, 0.04, 0.4, 2.0 ng/ml) or TNF (0, 0.4, 4.0 and 20 ng/ml) and control, SCF (10 ng/ml) or IL-3 (10 ng/ml). Data presented are the mean and SE.

Figure 2: Time course of HIV expression by ceramide treated cells.

Panel A: HIV-1 core p24 production by U-1_{MB} cells. ●: control, ■: ceramide treated cells.

Panel B: HIV-1 core p24 production by OM-10.1 cells treated with 50 μ M ceramide. ●: control, ■: ceramide treated cells.

Panel C: Relative cell growth in the absence or presence of 50 μ M ceramide. ▲ and Δ: OM-10.1; ■ and □: U-1_{MB} cells. Open symbols represent controls, full symbols represent ceramide treated cells. Initial cell density: 5×10^5 cells/ml.

Panel D: Cell viability in the absence (open symbols) or presence of 50 μ M ceramide.

Figure 3: Enhancement of CAT transcription driven by the HIV LTR by ceramide. U-937 cells transfected by the pU3R-III CAT/h plasmid were exposed to increasing concentrations of ceramide, DMSO vehicle or PMA as a positive control and CAT activities were determined at day 5. Ceramide induced a dose dependent increase in CAT activity.

Figure 4: Interaction of the nuclear receptors ARP-1, EAR-3, EAR-2, HNF-4, NGFI-B, and RXR- α with sequences located within the HIV-1 LTR.

- Panel A:** Schematic representation of the HIV-1 LTR. Nucleotide numbers are relative to the transcription start site (+1). The double-stranded oligo NRRE-1 spanning the -320 to -358 LTR region is bracketed. Asterisks and open circles indicate nucleotides whose methylation interfered strongly and weakly, respectively, with receptor binding. The coding strand of the double-stranded oligo NMUT-BD harboring eight nucleotide substitutions (white letters) is also shown. NRRE-1 and NMUT-BD oligos have a 5' GATC overhang.
- Panel B:** DNase I protection of the -453 to -244 DNA region of the HIV_{LAI} LTR ³²P-labeled at the coding or noncoding strands in the presence of COS-1-produced receptors, as indicated. Lanes G+A: chemical sequencing ladder; control: DNase I protection using whole-cell extracts from mock-transfected COS-1 cells. Solid bars indicate areas protected by ARP-1, EAR-3, and EAR-2; open bars indicate areas protected by HNF-4.
- Panel C:** Methylation interference analysis of in vitro translated ARP-1, EAR-3, EAR-2, and HNF-4 with an oligo probe spanning the -368 to -310 region of the HIV_{LAI} LTR ³²P-labeled at the coding or noncoding strands, as indicated. Lanes M: G+A sequencing ladder; B: protein-bound probe; and F: free probe. Methylation interference symbols as in A.
- Panel D:** Electrophoretic mobility shift analysis or nuclear receptor binding to the NRRE-1 probe in the presence (+) or absence (-) of 100-fold molar excess of the indicated unlabeled oligo

Final Report
(Period 09/28/90 - 09/27/93)
Contract No: DAMD17-90-C-0106

competitors and 9-cis retinoic acid (10^{-6} M).
NS: non-specific oligo with unrelated
sequence.

Figure 5: Retinoic acid effects on HIV-1 IIIB acute infection in T cell and monocytoïd cell lines. On day 17 after infection, p24 antigen level and MTT dye conversion was measured. To compensate for nonspecific effects on cell viability the ratio of p24 antigen/MTT dye conversion is presented.

Panel A: H9 and CEM cells

Panel B: U937 and THP-1 cells


Panel C: The same experiment was performed on THP-1 cells 3 months after infection

The data presented is the mean of three experiments, each performed in duplicate.





IC50

	ZDV uM	GAN uM	IFN U/ml	TNF ng/ml	TGF ng/ml
BFU CONT	0.5 +/- .1	.55 +/- .1	3500 +/- 1300	10 +/- 1	1.3 +/- .3
BFU SCF	0.8 +/- .2	.8 +/- .1	5000 +/- 500	18 +/- 10	1.3 +/- .8
BFU IL3	*1.0 +/- .2	.9 +/- .6	6000 +/- 1000	16 +/- 3	2.4 +/- .3
GM CONT	1.2 +/- .05	1.2 +/- .6	4000 +/- 400	25 +/- 13	1.5 +/- .4
GM SCF	1.1 +/- .1	1.7 +/- .3	5600 +/- 400	40 +/- 8	2.5 +/- 1.5
GM IL3	1.2 +/- .1	1.2 +/- .4	5200 +/- 400	25 +/- 5	2.5 +/- 1.3


(Table 1)






(Figure 1)




(Figure 2)



(Figure 3)



(Figure 4)



(Figure 5)